

Math 214 Final Exam

Practice Problems

Instructor: Jay Daigle

This is not a practice test, in the sense that it is not the format I expect the test to be. It is a collection of practice problems. I will update you when I finalize the test format.

Proofs

1. Let Q be the subspace of $\mathcal{P}(x)$ consisting of polynomials with zero constant term. Prove that the function $D : Q \rightarrow \mathcal{P}(x)$ given by the derivative is an isomorphism.
2. Let $U = \text{span}\{x, \sin(x), \cos(x), x^5, 1\}$. Find an isomorphism between U and \mathbb{R}^5 .
3. Suppose V is a vector space and $L : V \rightarrow \mathbb{R}^5$ is surjective and $\dim \ker(L) = 2$. What can you say about V ?
4. Suppose $T : \mathbb{R}^5 \rightarrow \mathcal{P}_4(x)$ and $\dim \ker(T) = 1$. What can you say about $T(\mathbb{R}^5)$?
5. If λ is an eigenvalue of A then prove that λ^{-1} is an eigenvalue of A^{-1} .

Things to Ponder

1. Find a 4×4 matrix with no real eigenvalues. Is it possible to find a 3×3 matrix with no real eigenvalues?
2. Find matrices $A, B \in M_n \times M_n$ such that $\text{Tr}(A)\text{Tr}(B) \neq \text{Tr}(AB)$.
Find a matrix A such that $\text{Tr}(A^2) < 0$.
3. What happens if you use the Gram-Schmidt process on a set of vectors that isn't linearly independent?

Find the transition matrices between the following bases

1. The standard basis and

$$F = \left\{ \begin{bmatrix} 5 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 3 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ 6 \\ 3 \end{bmatrix} \right\}$$

2. The standard basis and

$$F = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right\}$$

- 3.

$$E = \left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} \right\} \quad \text{and} \quad F = \left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \right\}$$

4.

$$E = \left\{ \begin{bmatrix} 3 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 3 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \right\} \quad \text{and} \quad F = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$$

Write the given element in the given basis

1. Write $(3, 1, 4)$ in the basis $F = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$.
2. Write $(2, 7, 1)$ in the basis $F = \left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix} \right\}$.
3. Write $(1, -1, 0)$ in the basis $F = \left\{ \begin{bmatrix} 3 \\ 5 \\ 2 \end{bmatrix}, \begin{bmatrix} 7 \\ 1 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$.
4. Write $(2, 3, 4)$ in the basis $F = \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \right\}$.

Find the matrix of the operator with respect to the given basis

1. Give the matrix of $L(x, y, z) = (3x + y + z, 5x - 2y + z, y + z)$ with respect to $F = \left\{ \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix} \right\}$.
2. Give the matrix of $L(x, y, z) = (2x + 3y - z, 4x - y + 3z, 2x + z)$ with respect to $F = \left\{ \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \right\}$.
3. Give the matrix of $L(x, y, z) = (-x + 4y + 2z, 3x - 5y + 2, 3x + 2y)$ with respect to $F = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\}$.
4. Give the matrix of $L(x, y, z) = (2x - y, 3x + y + 4z, x + 2y + z)$ with respect to $F = \left\{ \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\}$.

Angles and Magnitudes

1. Compute

$$\begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 5 \\ 7 \\ -1 \end{bmatrix}, \quad \begin{bmatrix} 4 \\ 1 \\ 3 \\ 5 \end{bmatrix} \cdot \begin{bmatrix} 2 \\ -5 \\ 7 \\ 4 \end{bmatrix}, \quad \begin{bmatrix} 1 \\ 3 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 4 \\ -1 \\ -1 \end{bmatrix}, \quad \begin{bmatrix} 7 \\ 1 \\ 5 \end{bmatrix} \cdot \begin{bmatrix} -3 \\ 1 \\ 1 \end{bmatrix}.$$

2. Find the magnitudes and corresponding unit vectors for

$$\begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}, \quad \begin{bmatrix} 5 \\ 12 \end{bmatrix}, \quad \begin{bmatrix} 4 \\ 2 \\ -2 \end{bmatrix}, \quad \begin{bmatrix} 7 \\ -1 \\ -3 \end{bmatrix}.$$

3. Find $\text{proj}_{\mathbf{v}} \mathbf{u}$ for

(a) $\mathbf{u} = (5, 2), \mathbf{v} = (-3, 4)$

(b) $\mathbf{u} = (2, 1), \mathbf{v} = (7, 1)$

(c) $\mathbf{u} = (3, 1, 4), \mathbf{v} = (2, 1, 1)$

(d) $\mathbf{u} = (2, 1, 1), \mathbf{v} = (-4, -1, -1)$

(e) $\mathbf{u} = (5, 0, 0), \mathbf{v} = (3, 2, 1)$.

Diagonalization Theory

1. In class we saw that

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -1 & 3 & -1 \\ 1 & -2 & 1 \\ 1 & -3 & 2 \end{bmatrix} \begin{bmatrix} 2 & -3 & 1 \\ 1 & -2 & 1 \\ 1 & -3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 3 & -1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}.$$

Multiply out the three matrices on the right and confirm that this works.

2. Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$. What are the eigenvalues of A ? Is $A^2 = A$? Why not?

3. Show the following pairs of matrices are not similar:

$$A = \begin{bmatrix} 4 & 1 \\ 3 & 1 \end{bmatrix}$$

$$B = \begin{bmatrix} 1 & 5 \\ 1 & 1 \end{bmatrix}$$

$$C = \begin{bmatrix} 2 & 1 & 4 \\ 0 & 2 & 3 \\ 0 & 0 & 4 \end{bmatrix}$$

$$D = \begin{bmatrix} 3 & 0 & 0 \\ 2 & 2 & 0 \\ 5 & 1 & 3 \end{bmatrix}$$

$$E = \begin{bmatrix} 3 & 4 & 1 \\ 0 & 8 & -2 \\ 0 & 0 & 10 \end{bmatrix}$$

$$F = \begin{bmatrix} 4 & 0 & 0 \\ -1 & 5 & 0 \\ 5 & 3 & 12 \end{bmatrix}$$

$$G = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$H = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Diagonalization

For each of the following matrices, determine whether it is diagonal. If it is, diagonalize it, then compute A^5 .

1. $A = \begin{bmatrix} 5 & 2 \\ 2 & 5 \end{bmatrix}$

2. $A = \begin{bmatrix} -4 & 6 \\ -3 & 5 \end{bmatrix}$

3. $A = \begin{bmatrix} 3 & 1 & 0 \\ 0 & 3 & 1 \\ 0 & 0 & 3 \end{bmatrix}$

4. $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$

$$5. A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 2 & 1 \\ 3 & 0 & 1 \end{bmatrix}$$

$$6. A = \begin{bmatrix} 2 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 0 & 2 \end{bmatrix}$$

Orthogonality and Projection

- Suppose $\|\mathbf{u}\| = 3$, $\|\mathbf{u} + \mathbf{v}\| = 4$, $\|\mathbf{u} - \mathbf{v}\| = 6$. Find $\|\mathbf{v}\|$.
- Find the orthogonal complement (in \mathbb{R}^n) of the following spaces:

$$W = \{(2t, -t) : t \in \mathbb{R}\}$$

$$W = \text{span}\{(2, -1, 3)\}$$

$$W = \{(t, -t, 3t) : t \in \mathbb{R}\}$$

$$W = \text{span}\{(1, -1, 3, -2), (0, 1, -2, 1)\}.$$

- Find the orthogonal decomposition of
 - $(7, -4)$ with respect to $\text{span}\{(1, 1)\}$
 - $(1, 2, 3)$ with respect to $\text{span}\{(2, -2, 1), (-1, 1, 4)\}$
 - $(4, -2, 3)$ with respect to $\text{span}\{(1, 2, 1), (1, -1, 1)\}$
 - $(3, 2, -3, 4)$ with respect to $\text{span}\{(2, 1, 0, 1), (0, -1, 1, 1)\}$.
 - $(2, -1, 5, 6)$ with respect to $U = \text{span}\{(1, 1, 1, 0), (1, 0, -1, 1)\}$.
- Let $V = \mathcal{P}_2(x)$ and define $\langle f, g \rangle = f(-1)g(-1) + f(0)g(0) + f(1)g(1)$.
 - Find the projection of $3x - 4x^2$ onto the vector $1 + x + x^2$.
 - Find the orthogonal decomposition of $2 + x$ with respect to the spaces $W = \text{span}\{5 + x\}$ and $W^\perp = \text{span}\{2 - 3x^2, -2 + 5x + 2x^2\}$. (You can assume that the space I gave you is in fact W^\perp . But you can also check yourself, for practice.)
 - Find the orthogonal decomposition of $3 - 3x + x^2$ with respect to $W = \{3 - 5x, 4x - 3x^2\}$ and $W^\perp = \{2 + 3x + 2x^2\}$.
 - Find the orthogonal complement of $W = \{\alpha_0 + \alpha_2 x^2 : \alpha_0, \alpha_2 \in \mathbb{R}\}$.