Math 1231-13: Single-Variable Calculus 1 George Washington University Spring 2024 Recitation 6

Jay Daigle

Friday February 23, 2024

Problem 1. Suppose a particle has height as a function of time given by $h(ts) = (2t^3 - 3t^2 - 12t + 3)$ m.

- (a) What is the velocity of this particle at time t = 0? What are the units, and why?
- (b) What is the acceleration of this particle at time t = 0? What are the units and why?
- (c) When is the particle speeding up? When is it slowing down?

Problem 2. Suppose that p(t) = 10 - 2t is momentum (in kg m/s) of a ball thrown directly upwards, as a function of time (in seconds).

- (a) What units does the derivative p'(t) take as input? What units are its output? (Do you know of any physical quantity that's represented by those units?)
- (b) What does the derivative p'(t) represent physically? What would it mean for p'(t) to be big, or small?
- (c) Calculate p'(3). What does this tell you? What physical observation could you measure to check if your calculation was correct?

Problem 3. Suppose the cost of buying *m* machines is $C(m) = 500 + 10m + .05m^2$. There's some start-up cost to having any machines at all; then each machine costs a bit more than the previous one.

- (a) What are the units of the inputs to the function C? What are the units of the outputs?
- (b) What is C(1)? C(10)? C(100)?

- (c) Find a formula for C'(m). What are the units of the input and output to C'(m)?
- (d) What is C'(10)? How should we interpret this number?
- (e) What is the *average* cost per machine when you have ten machines? How does this compare to your previous answer?
- (f) What is C''(m)? What are the units? What is C''(10) and how should we interpret it?

Problem 4 (Bonus). Let Q(p) = 10000 - 10p give the number of widgets you can sell at a given price p.

- (a) If you set a price of \$100, how many widgets will you be able to sell? What if you set a price of \$1000?
- (b) What is the derivative of Q? What are its units?
- (c) What is Q'(100) and what does that tell you?

Problem 5. Find an equation for the tangent line to $y = 6 \cos x$ at $(\pi/3, 3)$.

Problem 6. Find an equation for the tangent line to $y \cos(x) = 1 + \sin(xy)$ at the point (0, 1).

Problem 7. Suppose we have some function f such that $8f(x) + x^2(f(x))^3 = 24$, and we know that f(4) = 1. (Say we've measured this experimentally and now want to understand or compute with the function). Now suppose we want to estimate f(5) (without having to solve the equation).

- (a) Use implicit differentiation on the equation $8f(x) + x^2(f(x))^3 = 24$ to find a formula relating x, f(x) and f'(x).
- (b) Use this formula to determine f'(4).
- (c) Find a formula for the linear approximation of f near 4.
- (d) Estimate f(5).

Problem 8 (Bonus). (a) If $\sqrt{xy} = x^2y - 2$, find a formula for $\frac{dy}{dx}$ in terms of x and y.

(b) Find an equation of the tangent line at the point (1, 4).