Math 1231 Practice Midterm Solutions

Instructor: Jay Daigle

Problem 1 (M3). (a) Find and classify all the critical points of $f(x) = (x-5)\sqrt[3]{x^2}$. [Note: this is quite hard but it's good practice.]

Solution: We compute

$$f'(x) = \sqrt[3]{x^2} + (x-5)\frac{2}{3}x^{-1/3} = x^{2/3} + \frac{2x-10}{3\sqrt[3]{x}}$$
$$= \frac{3x+2x-10}{3\sqrt[3]{x}} = 5\frac{x-2}{3\sqrt[3]{x}}$$

This is equal to zero when $x = 2$ and is undefined when $x = 0$, so the two critical points are $x = 0$ and $x = 2$.

We could try to use the second derivative test here, but it won't really work. We get

$$f''(x) = \frac{10x+10}{9x^{4/3}}$$
$$f''(2) = \frac{5}{3\sqrt[3]{2}} > 0$$

so we see that f has a local minimum at $x = 2$, but $f''(0)$ is undefined so it doesn't help us classify $x = 0$.

Instead we compute a chart

	$5(x-2)$	$\frac{1}{3\sqrt[3]{x}}$	$f'(x)$
$x < 0$	$-$	$-$	$+$
$0 < x < 2$	$-$	$+$	$-$
$2 < x$	$+$	$+$	$+$

Thus we conclude that f has a local maximum at $x = 0$ and a local minimum at $x = 2$.

(b) The function $g(x) = (x^2 - 3x)\sqrt[3]{x-3}$ has absolute extrema either on $(-4,-1)$ or on $[1,4]$. Pick one of those intervals, explain why g has extrema on that interval, and find the absolute extrema.

Solution: We compute

$$g'(x) = (2x-3)\sqrt[3]{x-3} + (x^2-3x)\frac{1}{3}(x-3)^{-2/3}.$$

This is undefined when $x = 3$, and to find the zeroes we compute

$$0 = (2x-3)\sqrt[3]{x-3} + (x^2-3x)\frac{1}{3}(x-3)^{-2/3}$$
$$= (2x-3)\sqrt[3]{x-3} + \frac{x^2-3x}{3(x-3)^{2/3}}$$
$$0 = (2x-3)\sqrt[3]{x-3}\cdot 3(x-3)^{2/3} + (x^2-3x)$$
$$= (2x-3)3(x-3) + (x^2-3x) = 3(2x^2-3x-6x+9) + x^2 - 3x$$
$$= 7x^2 - 30x + 27 = (7x-9)(x-3)$$

which is zero when $x = 3$ or $x = 9/7$. So then we have

$$\begin{aligned} g(1) &= -2\sqrt[3]{-2} = 2\sqrt[3]{2} \\ g(9/7) &= \left(\frac{81}{49} - \frac{27}{7}\right)\sqrt[3]{-12/7} = \frac{108}{49}\sqrt[3]{12/7} \\ g(3) &= 0 \\ g(4) &= 4 \cdot \sqrt[3]{1} = 4. \end{aligned}$$

All of these numbers are non-negative, so the minimum value is 0, at $x = 3$. We can see that $2\sqrt[3]{2} < 2 \cdot 2 = 4$, and maybe convince ourselves that

$$\frac{108}{49}\sqrt[3]{12/7} \approx 2 \cdot 1 < 4.$$

Thus the maximum value is 4, which occurs at 4.

Problem 2 (S5)**.** Find a tangent line to the curve given by $x^4 - 2x^2y^2 + y^4 = 16$ at the point $(\sqrt{5}, 1)$.

Solution: We use implicit differentiation, and find that

$$\begin{aligned} 4x^3 - 2\left((2xy^2 + x^2 2y\frac{dy}{dx}\right) + 4y^3\frac{dy}{dx} &= 0 \\ 4x^3 - 4xy^2 &= 4x^2y\frac{dy}{dx} - 4y^3\frac{dy}{dx} \\ \frac{4x^3 - 4xy^2}{4x^2y - 4y^3} &= \frac{dy}{dx} \end{aligned}$$

Thus at the point $(\sqrt{5}, 1)$ we have

$$\frac{dy}{dx} = \frac{4\sqrt{5}^3 - 4\sqrt{5}\cdot 1^2}{4\sqrt{5}^2 \cdot 1 - 4 \cdot 1^3} = \sqrt{5}\left(\frac{20-4}{20-4}\right) = \sqrt{5}.$$

Thus the equation of our tangent line is

$$\begin{aligned} y - y_0 &= m(x - x_0) \\ y - 1 &= \sqrt{5}(x - \sqrt{5}). \end{aligned}$$

Problem 3 (S6)**.** Consider this baseball diamond, which is a square with 90ft sides. A batter hits the ball and runs from home toward first base at a speed of 22ft/s. At what rate is the distance between the runner and second base changing when the runner has run 30ft?



Solution: We use the Pythagorean theorem, $a^2 + b^2 = c^2$, where a is the distance of the runner from first base and b is the distance of second base from first base. Then c is the distance between the runner and second base, which we want to know about, and we have it related to a and b, which we do know about.

When the runner has run 30ft, then we have $a = 60$ft and $b = 90$ft is a constant. Then we have

$$\begin{aligned} c^2 = a^2 + b^2 = 60^2 + 90^2 = 3600 + 8100 = 11700 \\ c = 10\sqrt{117} = 30\sqrt{13}. \end{aligned}$$

Alternatively,

$$\begin{aligned} c^2 = 60^2 + 90^2 = 30^2(2^2 + 3^2) \\ c = 30\sqrt{13}. \end{aligned}$$

We know that $a' = -22$ft/s, so we compute

$$\begin{aligned} 2aa' + 2bb' &= 2cc' \\ aa' + bb' &= cc' \\ 60\text{ft} \cdot (-22)\text{ft/s} + 90\text{ft} \cdot 0\text{ft/s} &= 30\sqrt{13}\text{ft} \cdot c' \\ 2 \cdot -22\sqrt{13}\text{ft/s} &= c'. \end{aligned}$$

So the distance between the runner and second base is decreasing at $\frac{44}{\sqrt{13}} \approx 12.2$ feet per second.

Problem 4 (S7)**.** Let $f(x) = \dfrac{x^3 - 2}{x^4}$. We compute that $f'(x) = \dfrac{8 - x^3}{x^5}$ and $f''(x) = \dfrac{2x^3 - 40}{x^6}$. Sketch a graph of f.

Your answer should discuss the domain, asymptotes, roots, limits at infinity, critical points and values, intervals of increase and decrease, and concavity.

Solution: The function is defined everywhere except at 0. Near zero, we can see the top is always negative and the bottom is always positive, so $\lim_{x\to 0} f(x) = -\infty$ and we should have a downwards asymptote on either side.

We see there is a root at $x = \sqrt[3]{2}$, and $\lim_{x\to\pm\infty} f(x) = 0$.

We see that $f'(x)$ is undefined at $x = 0$, and is zero when $x^3 = 8$ and thus when $x = 2$. So our critical points occur at 0 and 2. We calculate $f(2) = \frac{6}{16}$, and f isn't defined at 0. By making a chart, we get

	$8 - x^3$	x^5	$f'(x)$
$x < 0$	$+$	$-$	$-$
$0 < x < 2$	$+$	$+$	$+$
$2 < x$	$-$	$+$	$-$

so f is decreasing for values less than zero or greater than 2, and increasing for values between 0 and 2.

The second derivative is undefined at 0, and is zero when $2x^3 - 40$ and so when $x = \sqrt[3]{20}$, so our potential points of inflection are 0, $\sqrt[3]{20}$. We compute $f(\sqrt[3]{20}) = \frac{18}{20\sqrt[3]{20}}$. We can make a chart again, but we see that the denominator of $f''(x) \geq 0$, so $f''(x) > 0$ if $x > \sqrt[3]{20}$ and $f''(x) < 0$ if $x < \sqrt[3]{20}$.



Problem 5 (S8)**.** We want to build a rectangular fence that will enclose 200m^2. One pair of parallel sides cost \$3/m and the other pair costs \$8/m. What dimensions minimize the cost of the fence? Justify your claim that this is a minimum.



Solution: We want to minimize $3w + 8\ell$ subject to the constraint $\ell w = 200$. Thus we have $w = 200/\ell$, and then our function is $C(\ell) = 600/\ell + 8\ell$. We get

$$\begin{aligned} C' &= -600/\ell^2 + 8 = 0 \\ -8\ell^2 &= -600 \\ \ell^2 &= 75 \\ \ell &= 5\sqrt{3}. \end{aligned}$$

Then we have $\ell = 5\sqrt{3}$ and $w = \frac{200}{5\sqrt{3}} = \frac{40}{3}\sqrt{3}$.

We have two options for proving this is a maximum (we only need one):

(a) Extreme Value Theorem: We can't really use the EVT here because we don't have a closed interval.

(b) First Derivative Test: For $\ell < 5\sqrt{3}$ we have $C' < 0$ so the function is decreasing, and for $\ell > 5\sqrt{3}$ we have $C' <>$ so the function is increasing. Thus we have a unique minimum at $5\sqrt{3}$.

(c) Second derivative test: $C''(\ell) = -1200/\ell^3$. Then $C''(\ell) > 0$ for ℓ?0, which implies there is a single relative minimum, at $\ell = 5\sqrt{3}$. This doesn't really rigorously prove that this is an absolute maximum but I'll take it.